

A REFINEMENT OF THE HOFER– ZEHNDER THEOREM ON THE EXISTENCE OF CLOSED TRAJECTORIES NEAR A HYPERSURFACE

LEONARDO MACARINI¹ AND FELIX SCHLENK²

ABSTRACT. The Hofer–Zehnder theorem states that almost every hypersurface in a thickening of a hypersurface S in a symplectic manifold (M, ω) carries a closed characteristic provided that S bounds a compact submanifold and (M, ω) has finite capacity. We show that it is enough to assume that the thickening of S has finite capacity.

We consider a smooth symplectic manifold (M, ω) . A *hypersurface* S in M is a smooth compact connected orientable codimension 1 submanifold of $M \setminus \partial M$ without boundary. A characteristic on S is an embedded circle in S all of whose tangent lines belong to the distinguished line bundle

$$\mathcal{L}_S = \{(x, \xi) \in TS \mid \omega(\xi, \eta) = 0 \text{ for all } \eta \in T_x S\}.$$

We denote by $\mathcal{P}(S)$ the set of closed characteristics on S . Examples show that $\mathcal{P}(S)$ can be empty, see [2, 3]. We therefore follow [4] and consider parametrized neighbourhoods of S . Since S is orientable, there exists an open neighbourhood I of 0 and a smooth diffeomorphism

$$\psi: S \times I \rightarrow U \subset M$$

such that $\psi(x, 0) = x$ for $x \in S$. We call ψ a *thickening* of S , and we abbreviate $S_\epsilon = \psi(S \times \{\epsilon\})$.

Given an open subset $U \subset M$ we consider the function space $\mathcal{H}(U)$ of smooth functions $H: U \rightarrow [0, \max H]$ such that

- $H|_V = 0$ for some nonempty open set $V \subset U$;
- $H|_{U \setminus K} = \max H$ for some compact set $K \subset U$.

We say that $H \in \mathcal{H}(U)$ is *HZ-admissible* if the flow φ_H^t has no non-constant T -periodic orbit with period $T \leq 1$, and we set

$$\mathcal{H}_{\text{HZ}}(U) = \{H \in \mathcal{H}(U) \mid H \text{ is HZ-admissible}\}.$$

Date: February 1, 2008.

¹ Partially supported by CNPq-PROFIX, Brazil.

² Supported by the von Roll Research Foundation.

The Hofer–Zehnder capacity of U is defined as

$$c_{\text{HZ}}(U) = \sup \{ \max H \mid H \in \mathcal{H}_{\text{HZ}}(U) \}.$$

It has been shown in [4, Sections 4.1 and 4.2] that for any thickening $\psi: S \times I \rightarrow U \subset M$ for which $c_{\text{HZ}}(U) < \infty$, the set $\{\epsilon \in I \mid \mathcal{P}(S_\epsilon) \neq \emptyset\}$ is dense in I , and that

$$\mu \{ \epsilon \in I \mid \mathcal{P}(S_\epsilon) \neq \emptyset \} = \mu(I)$$

if S bounds a compact submanifold of M and $c_{\text{HZ}}(M) < \infty$. Here, μ denotes Lebesgue measure. In this note we prove

Theorem 1. *For any thickening $\psi: S \times I \rightarrow U \subset M$ we have*

$$\mu \{ \epsilon \in I \mid \mathcal{P}(S_\epsilon) \neq \emptyset \} = \mu(I)$$

provided that $c_{\text{HZ}}(U) < \infty$.

Proof. Consider a thickening $\psi: S \times I \rightarrow U \subset M$. We can assume that $I =]-1, 1[$. Let $G = \{\epsilon \in]-1, 1[\mid \mathcal{P}(S_\epsilon) \neq \emptyset\}$ be the parameters of the good hypersurfaces.

Step 1. *The set G is measurable.*

Proof. We define the smooth function $K: U \rightarrow]-1, 1[$ by

$$K(z) = \epsilon \quad \text{if } z \in S_\epsilon.$$

The set $\mathcal{P}_K(\epsilon)$ of closed orbits of the Hamiltonian flow φ_K^t of K on S_ϵ corresponds to $\mathcal{P}(S_\epsilon)$. For each closed orbit x of φ_K^t we denote by $T(x)$ its period defined as

$$T(x) = \min \{ t > 0 \mid \varphi_K^t(p) = p \}$$

where p is any point on x . For each $n \in \mathbb{N}$ we set

$$G_n = \{ \epsilon \in]-1, 1[\mid \text{there exists } x \in \mathcal{P}_K(\epsilon) \text{ such that } T(x) \leq n \}.$$

It follows from the Arzelà–Ascoli Theorem that G_n is a closed subset of $]-1, 1[$, see [4, page 109, Proposition 1] The set $G = \bigcup_{n \geq 1} G_n$ is thus measurable.

Step 2. *The set G has full measure.*


Proof. Applying the argument in the proof of Theorem 3 in Section 4.2 of [4] to Hamiltonian functions $H: U \rightarrow \mathbb{R}$ as in the figure below such that

$$H(x) = f(|K(x)|) \quad \text{if } x \in S_{-\epsilon} \cup S_\epsilon, \quad \epsilon \leq t \leq 1,$$

we see that $\mathcal{P}(S_{-\epsilon}) \neq \emptyset$ or $\mathcal{P}(S_\epsilon) \neq \emptyset$ whenever the function $C(t): [0, 1[\rightarrow \mathbb{R}$, $t \mapsto c_{\text{HZ}}(\psi(S \times]-t, t])$ is Lipschitz continuous at ϵ . Since $C(t)$ is

monotone increasing, it is differentiable almost everywhere, and thus Lipschitz continuous almost everywhere. Therefore,

$$(1) \quad \mu \{ \epsilon \in [0, 1[\mid \mathcal{P}(S_{-\epsilon}) \neq \emptyset \text{ or } \mathcal{P}(S_{\epsilon}) \neq \emptyset \} = 1.$$



Let $B =]-1, 1[\setminus G$ be the parameters of the bad hypersurfaces. Arguing by contradiction we assume that $\mu(B) > 0$. Choose an interval $]a, b[\subset]-1, 1[$ such that

$$(2) \quad \mu(]a, b[\cap B) \geq \frac{2}{3}(b - a).$$

Applying (1) to $]a, b[$ instead of $] - 1, 1[$ we see that

$$(3) \quad \mu(]a, b[\cap G) \geq \frac{1}{2}(b - a).$$

Combining (2) with (3) we find that

$$\begin{aligned} b - a &= \mu(]a, b[) \\ &= \mu(]a, b[\cap B) + \mu(]a, b[\cap G) \\ &\geq \frac{2}{3}(b - a) + \frac{1}{2}(b - a) \\ &> b - a. \end{aligned}$$

This contradiction shows that $\mu(B) = 0$, and so $\mu(G) = \mu(]-1, 1[)$. $\square$

It is often of interest to find closed characteristics in a restricted set of homotopy classes. We thus fix a subset Γ of the fundamental group $\pi_1(S)$, and given a thickening $\psi: S \times I \rightarrow U \subset M$, we denote by $\mathcal{P}^\Gamma(S_\epsilon)$ the set of closed characteristics on S_ϵ representing an element of $\Gamma \subset \pi_1(S_\epsilon) = \pi_1(S)$. We can assume that $I =]-1, 1[$, and for each $\epsilon \in]0, 1[$ we set $U_\epsilon = \psi(S \times]-\epsilon, \epsilon[)$. We say that $H \in \mathcal{H}(U_\epsilon)$ is HZ^Γ -admissible if the flow φ_H^t has no non-constant T -periodic orbit with period $T \leq 1$ which represents an element of $\Gamma \subset \pi_1(U_\epsilon) = \pi_1(S)$, and we set

$$\mathcal{H}_{\text{HZ}}^\Gamma(U_\epsilon) = \{H \in \mathcal{H}(U_\epsilon) \mid H \text{ is } \text{HZ}^\Gamma\text{-admissible}\}.$$

The Γ -sensitive Hofer-Zehnder capacity of U_ϵ is defined as

$$c_{\text{HZ}}^\Gamma(U_\epsilon) = \sup \{ \max H \mid H \in \mathcal{H}_{\text{HZ}}^\Gamma(U_\epsilon) \}.$$

Notice that the map $\epsilon \mapsto c_{\text{HZ}}^\Gamma(U_\epsilon)$ is monotone increasing. Repeating the above proof with c_{HZ} replaced by c_{HZ}^Γ we find the following refinement of Theorem 1.

Theorem 2. *For any thickening $\psi: S \times I \rightarrow U \subset M$ of a hypersurface S in (M, ω) we have*

$$\mu \{ \epsilon \in I \mid \mathcal{P}^\Gamma(S_\epsilon) \neq \emptyset \} = \mu(I)$$

provided that $c_{\text{HZ}}^\Gamma(U) < \infty$.

While a version of Theorem 1 is applied in [1], Theorem 2 is used in [5] to show that the flow describing the dynamics of a charge in a rational symplectic magnetic field has a closed orbit on almost every small energy level.

Acknowledgements. This note was written during the second authors stay at Stony Brook in April 2003. He wishes to thank Ely Kerman and Dusa McDuff for their warm hospitality.

REFERENCES

- [1] U. Frauenfelder and F. Schlenk. Hamiltonian dynamics on convex symplectic manifolds. math.SG/0303282.
- [2] V. Ginzburg. Hamiltonian dynamical systems without periodic orbits. Northern California Symplectic Geometry Seminar, 35–48, *Amer. Math. Soc. Transl. Ser. 2*, **196**, Amer. Math. Soc., Providence, RI, 1999.
- [3] V. Ginzburg and B. Gürel. Relative Hofer–Zehnder capacity and periodic orbits in twisted cotangent bundles. math.DG/0301073.
- [4] H. Hofer and E. Zehnder. *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser, Basel, 1994.
- [5] L. Macarini. Hofer–Zehnder semicapacity of cotangent bundles and symplectic submanifolds. math.SG/0303230.

(L. MACARINI) INSTITUTO DE MATEMÁTICA PURA E APLICADA - IMPA,
ESTRADA DONA CASTORINA, 110 - JARDIM BOTÂNICO, 22460-320 RIO DE
JANEIRO RJ, BRAZIL

E-mail address: leonardo@impa.br

(F. SCHLENK) ETH ZÜRICH, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: schlenk@math.ethz.ch